




Contents lists available at ScienceDirect

Protein Expression and Purification

journal homepage: www.elsevier.com/locate/yprep



Cell-free expression of a functional pore-only sodium channel

Gabriela Kováčsová^{a,1}, Emil Gustavsson^a, Jiajun Wang^b, Mohamed Kreir^b, Sebastian Peuker^{a,*}, Sebastian Westenhoff^{a,*}

^a Department of Chemistry, University of Gothenburg, P.O. Box 462, SE-40530 Gothenburg, Sweden

^b Nanion Technologies GmbH, Gabrielenstraße 9, 80636 Munich, Germany

ARTICLE INFO

Article history:
Received 15 January 2015
and in revised form 1 March 2015
Available online xxx

Keywords:
Cell-free expression
Membrane protein
Sodium channel

ABSTRACT

Voltage-gated sodium channels participate in the propagation of action potentials in excitable cells. Eukaryotic Na_vs are pseudo homotetrameric polypeptides, comprising four repeats of six transmembrane segments (S1–S6). The first four segments form the voltage-sensing domain and S5 and S6 create the pore domain with the selectivity filter. Prokaryotic Na_vs resemble these characteristics, but are truly tetrameric. They can typically be efficiently synthesized in bacteria, but production *in vitro* with cell-free synthesis has not been demonstrated. Here we report the cell-free expression and purification of a prokaryotic tetrameric pore-only sodium channel. We produced milligram quantities of the functional channel protein as characterized by size-exclusion chromatography, infrared spectroscopy and electrophysiological recordings. Cell-free expression enables advanced site-directed labelling, post-translational modifications, and special solubilization schemes. This enables next-generation biophysical experiments to study the principle of sodium ion selectivity and transport in sodium channels.

© 2015 Published by Elsevier Inc.

Introduction

Voltage-gated sodium channels (Na_v)² are membrane proteins from the superfamily of voltage-gated ion channels, closely related to voltage-gated potassium channels and voltage-gated calcium channels [1]. Na_vs are present in all excitable cells, where they participate in the propagation of action potentials by changing the Na⁺ permeability of the cell membrane. Most voltage-gated ion channels comprise of similar building blocks and are mainly alpha-helical. Na_vs comprise six transmembrane segments (S1–S6), where S1–S4 form the voltage-sensing domain and S5 and S6 create the pore domain. In bacteria, four of these subunits arrange around a central pore to form a functional channel. In higher organisms, all four subunits are assembled from a single polypeptide chain, which may associate with auxiliary subunits [2].

Eukaryotic sodium channels were discovered first and have been the subject to extensive research for many decades [3].

However, the discovery of a prokaryotic bacterial sodium channel in 2001 [4] was a prerequisite for solving the proteins' three dimensional crystal structure [5], mainly because it enabled production of larger amounts of channel proteins. Bacterial Na_vs have 20–25% identity with human Na_vs and are expected to have a similar fold as they have nearly identical hydrophobicity profiles and predicted topologies in each of the pseudo-repeated eukaryotic domains [6]. Despite the available structural information, the mechanisms of molecular ion transport and ion selectivity are still not completely understood. X-ray crystallography catches high-resolution structures of stationary states, but lacks dynamic information. In principle, molecular dynamics simulations can be used to produce dynamical models. However, for potassium channels [7], such simulations have led to radically different proposals for mechanisms for ion transport [8,9]. This emphasizes the need for experimental validation. As we point out below, suitable experiments are today becoming possible. They may require sophisticated and site-selective modifications of the protein, which is asking for efficient production of Na_vs with cell-free (*in vitro*) synthesis.

In cell-free expression, proteins are expressed from exogenous template DNA added to the transcription and translation enzymes extracted from a cell lysate. This *in vitro* synthesis is becoming increasingly popular, particularly because it is possible to produce proteins which aggregate, or proteins which are toxic to host cells [10–14]. Today, typical yields of 0.3 mg to several milligrams of

* Corresponding authors at: Boehringer Ingelheim Pharma GmbH & Co. KG, Birkendorfer Straße 65, 88397 Biberach/Riß, Germany (S. Westenhoff).

E-mail addresses: peukers@googlemail.com (S. Peuker), westenho@chem.gu.se (S. Westenhoff).

¹ Current address: Department of Biomolecular Mechanisms, Max-Planck Institute for Medical Research, Jahnstraße 29, D-69120 Heidelberg, Germany.

² Abbreviations used: Na_vs, voltage-gated sodium channels; Na_vSp1p, sodium channel from *Silicibacter pomeroyi*; CMC, critical micelle concentration; GUVs, giant unilamellar vesicles.

protein per mL reaction mixture can be achieved in batch or in continuous mode, respectively, and popular reaction mixtures are extracted from *Escherichia coli* or wheat germs [14–16]. As cell-free expression gives direct access to the nascent polypeptide, it facilitates co-translational solubilization of membrane proteins in a wide range of detergents, lipids and nanodiscs [10,17]. Indeed, functional membrane proteins [10–13] including ion channels such as connexins [18], nicotinic acetylcholine receptors [19], drosophila olfactory receptors [20] and potassium channels can be produced by *in vitro* synthesis [17,21,22]. Also, eukaryotic sodium channels protein were synthesized *in vitro* [23]. Most of these examples were produced in cell extracts that yield microgram quantities of proteins, which is sufficient for electrophysiological studies, but not for spectroscopic and crystallographic investigations [18,19,21–23]. Isotope-labelled proteins are easily available as the amino acids are added to the reaction mixture as required [24].

In-vitro synthesis of ion channels is useful, because it may pave the way for the investigation of the molecular details of ion conductance in channel proteins. It has been demonstrated that *in vitro* synthesis enables time-saving direct reconstitution into oocytes [17]. Room temperature spectroscopy methods, such as NMR and vibrational spectroscopy, can in principle be used to study ion channels in native environments. Two-dimensional infrared spectroscopy is particularly interesting, because it can be used to characterize biological processes involving protein conformational change, e.g. transport or charge transfer with picosecond time resolution [25,26]. A specific infrared experiment to study the occupancy in selectivity filters of ion channels has been suggested [27]. Nonetheless, it is usually problematic to assign vibrational spectral signatures to specific sites in proteins, except for when these sites are labelled with isotopes or specific chemical groups [27–29]. *In-vitro* synthesis of protein makes possible site-directed labelling with specific amino acids using amber-stop codon technology [30,31]. To realize this new generation of spectroscopic experiments, reliable *in vitro* production of milligram quantities of ion channels is a prerequisite.

Here we report the production of a bacterial pore-only sodium channel from *Silicibacter pomeroyi* (Na_vSp1p) [32]. Designed and explored by the Minor group, the pore domain folds independently of the voltage-sensing domain into a functional channel protein, which displays selectivity for sodium over potassium ions. When produced in *E. coli*, it is more stable and expresses at higher levels than the complete channel [32]. We show here that cell-free production of a few milligrams of Na_vSp1p is possible, that the protein is folded correctly, and that a functional sodium channel is produced.

Material and methods

Extract cultivation, preparation

S12 extract was prepared from BL21 (DE3):RFI-CBD₃ [31] as described in [33]. Briefly, cells were grown in a 20 L fermentor (Braun Biostat C) at 37 °C in 2xYPTG medium supplemented with choline chloride (Fluka 28.6 mg/L), nicotinic acid (Acros, 25.1 mg/L), p-aminobenzoic acid (Aldrich, 20.0 mg/L), pantothenic acid calcium salt (Fluka, 9.4 mg/L), pyridoxal-5-phosphate (1.8 mg/L), (–)-riboflavin (3.9 mg/L), thiamine hydrochloride (USB Corporation, 17.7 mg/L), betaine hydrochloride (Calbiochem, 33.1 mg/L), D-biotin (MP Biomedicals, 0.1 mg/L), cyanocobalamin (Fluka, 0.01 mg/L), folic acid calcium salt hydrate (Sigma, 0.075 mg/L), iron (III) chloride hexahydrate (Scharlau Chemie S.A., 20.0 mg/L), sodium molybdate dehydrate (Acros, 3.5 mg/L), boric acid (1.2 mg/L), cobalt sulphate heptahydrate (4 mg/L), copper sulphate pentahydrate (Merck KGaA, 3.4 mg/L), manganese

sulphate hydrate (Merck KGaA, 1.9 mg/L), zinc sulphate heptahydrate (Scharlau Chemie S.A., 3.4 mg/L), and amino acids (Asp (28.5 mg/L), Gly (49.1 mg/L), His (9.35 mg/L), Ile (26.2 mg/L), Leu (29.9 mg/L), Lys (31.4 mg/L), Met (14.9 mg/L), Phe (15.3 mg/L), Pro (31.8 mg/L), Thr (37.7 mg/L), Trp (102.1 mg/L), Tyr (37.7 mg/L), Val (117.1 mg/L)). At OD₆₀₀ ~4.5 temperature was decreased to 10 °C by passing the cell suspension through a metal coil immersed in ice water, cells were harvested, washed with extraction buffer (10 mM Tris–acetate (pH 8.2), 14 mM Mg(OAc)₂, 60 mM K(OAc), cOMplete EDTA-free (Roche)), and finally resuspended in 10 mL extraction buffer/8 g of wet cells. The cells were lysed by a French press (two passages, 24,000 psi, ThermoFisher), centrifuged at 12,100g (10 min, 4 °C), the supernatant was decanted into fresh tubes and incubated for 2 h at a shaking incubator (30 °C, 150 rpm). We removed the release factor 1 protein from the cell extract. This is important for potential subsequent site-directed labelling steps with amber stop codons [30,34]. The S12 extract was passed over chitin resin (New England Biolabs) directly after the incubation and removal was confirmed by Western blotting. After addition of 1 mM DTT the S12 extract was dialyzed twice against extraction buffersupplemented with β-mercaptoethanol (1 mL/L), flash frozen, and stored at –80 °C.

Cell-free protein expression was performed in batch mode as described by [33]. Briefly, plasmid 0.01 µg/µL DNA, 14–20 mM Mg(OAc)₂, all 20 amino acids (1 mM each, besides Gln (4 mM) and Ser (2 mM)), 27.4 mM NH₄OH, 212 mM D-Glu, 230 mM KOH, 52.5 mM Hepes-KOH (pH 7.0), 1.1 mM ATP, 800 µM GTP, 800 µM CTP, 800 µM UTP, 640 µM cAMP, 68 µM folic acid (BioXtra), 1.7 mM DTT, 51.6 mM creatine phosphate (Roche), 4.4 mM L-(–)-malic acid (Fluka), 1.5 mM succinate (SAFC), 1.9 mM α-ketoglutaric acid (Fluka), 175 µg/mL tRNA (Roche), 8 U/mL RiboLock RNase Inhibitor (ThermoFischer), 1xComplete EDTA-free (Roche), 50 µg/mL T7RNA polymerase (prepared according to [35,36]), 125 µg/mL creatine kinase (Roche), 31% (v/v) S12, and detergents were mixed. Brij®-58 (polyoxyethylene (20) cetyl ether) and Brij®-78 (polyoxyethylene (18) octadecyl ether) (Sigma) were used at a final concentration of 10 times excess of the critical micelle concentration (CMC), 0.8 mM and 0.46 mM, respectively. DDM (*n*-dodecyl-β-D-maltoside) and DM (*n*-decyl-β-D-maltoside) (Anagrade Affymetrix) were used at 3 × CMC, 0.6 mM and 5.4 mM, respectively. Mixture was incubated for two hours at 30 °C and 800 rpm. For every batch of S12 the optimal Mg²⁺ was determined by GFPcyc3 expression. GFPcyc3 fluorescence was measured with a FluoStar plate reader (BMG Labtech, 390 nm (excitation), 520 nm (emission)).

To verify protein expression, Western blotting was performed following the manual for XCell II™ Blot Module and ONE-HOUR Western™ Basic Kit (Mouse) (GenScript) using Anti-His Antibody (GE Healthcare). Chemiluminescence was detected using a Fujifilm Las-1000 Luminescent Image Analyzer Chemi Fuji together with its software.

Overexpression of Na_vSp1p in *E. coli* cells

Na_vSp1p cloned into pHM3C-LIC, a vector containing a N-terminal hexa-His tag, a maltose-binding protein and a HRV 3C cleavage site was a kind gift from Daniel Minor [32]. The construct was transformed into *Escherichia coli* BL21 (DE3) and expression was performed as described [32].

Purification of Na_vSp1p from cell-free synthesis and expression in *E. coli*

We adapted a purification scheme from Shaya et al., 2011. After completed cell-free synthesis, the reaction mixture was centrifuged (16,000g, 20 min, 4 °C) and the supernatant was loaded onto a

gravity flow Ni-NTA agarose column (Qiagen) equilibrated with buffer A (20 mM Tris pH 8.0, 200 mM NaCl, 8% (vol/vol) glycerol, 2.7 mM DM). The column was washed with 7 column volumes of buffer A with 20 mM imidazole and eluted by 3 column volumes of buffer A with 300 mM imidazole. Large amounts, as e.g. channel protein produced in *E. coli* were loaded onto a His-Trap HP column using an Äcta system (GE Healthcare). The sample was desalted using buffer A either by dialysis (MWCO 12–14 kDa) or loaded onto a HiPrep 26/10 Desalting Column (GE Healthcare). The affinity tag and the maltose binding protein were cleaved by a His-labeled in-house HRV 3C protease (see below) at a ratio protein: protease = 1 mg: 0.28 mg at 8 °C overnight with gentle agitation. The protease was removed by a second Ni-NTA column which additionally removes traces of uncleaved protein and the His-tagged maltose binding protein tag. The last purification step was a size exclusion chromatography step (Superdex Increase 200 GL (GE Healthcare), with buffer C (20 mM Hepes pH 8.0, 200 mM NaCl, and 2.7 mM DM)). Protein purity was evaluated by SDS-PAGE stained with SimpleBlue™ SafeStain (life technologies) and the protein concentration was determined by absorbance at 280 nm using NanoDrop 1000 (Thermo Scientific), where the extinction coefficient for the uncleaved and cleaved protein was 90,300 M⁻¹ cm⁻¹ and 23,500 M⁻¹ cm⁻¹, respectively.

HRV 3C protease expression and purification

HRV 3C in a pET28 expression vector was obtained from Daniel Minor [32]. 10 mL 2YT (pH 7) containing kanamycin (100 µg/mL) was inoculated with a glycerol stock of pET28 in *E. coli* BL21 Star and grown overnight (220 rpm, 37 °C). Two liter cultures were inoculated with 6 mL of preculture and grown (180 rpm, 37 °C) to an O.D₆₀₀ of 0.6–0.8. Then the temperature was reduced to 22 °C and after 15 min, expression was induced with 0.4 mM IPTG (Affymetrix). After 19 h cells were harvested by centrifugation (6000g, 20 min, 4 °C). Pelleted cells were resuspended in 50 mL lysis buffer (50 mM Tris-HCl pH 8.0, 300 mM NaCl, 20 mM imidazole, 10% glycerol, 1 mg lysozyme from chicken egg white, Sigma and 1 mM PMSF) and disrupted using French Press, two passages at 24,000 psi. Cell lysate was separated from unbroken cells and cell debris by ultracentrifugation (42,000g, 1 h, 4 °C). The supernatant containing the octa-histidine tagged 3C protease was loaded onto a 5 mL HisTrap™ HP column (GE Healthcare) and equilibrated with chelating buffer (50 mM Tris-HCl pH 8.0, 300 mM NaCl, 20 mM imidazole, 10% glycerol). The column was washed with chelating buffer containing 46 mM imidazole and the bound protein was eluted by step application of elution buffer (50 mM Tris-HCl pH 8.0, 300 mM NaCl, 150 mM imidazole, 10% glycerol). The fractions containing the protein were collected and dialyzed overnight against 2 L dialysis buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 10 mM EDTA, 20% glycerol, 5 mM DTT) (MWCO 12–14 kDa, 8 °C). The concentration was measured photometrically using an extinction coefficient of 5960 M⁻¹ cm⁻¹ at 280 nm and M.W. was 21.282 kDa. Aliquots were flash frozen and stored at –20 °C.

IR spectroscopy

IR spectrum of the liquid sample was obtained with Agilent Cary 630 FTIR spectrometer with Diamond ATR Accessory. Buffer absorbance was recorded, scaled, and subtracted from sample spectrum as to minimize the water absorbance.

Functional characterization of reconstituted Na_vSp1p

Purified Na_vSp1p was incorporated into giant unilamellar vesicles (GUVs) using procedures described previously [32,37,38]. First,

GUVs were produced by electroformation using 10 mM 1,2, diphytanoyl-sn-glycero-3-phosphocholine and 1 mM cholesterol dissolved in trichloromethane. No phosphatidyl glycerol, phosphatidyl serine nor phosphatidylinositol were added as the lipid sensitivity of pore-only Na_vSp1p seems to be drastically reduced when compared to full length protein [39]. Approximately 20 µL of lipid solution was placed on the Vesicle Prep Pro (Nanion Technologies) ITO glass surface and air-dried. The dry lipid film was rehydrated using 250 µL 1 M sorbitol. GUVs were formed by electroswelling under the influence of an alternating electrical field for 2 h. GUVs were collected and incubated with protein suspension containing 0.5 µg/mL Na_vSp1p in 20 mM Hepes pH 8.0, 200 mM NaCl, and 2 mM DM for 15 min. Excess detergent was adsorbed by addition of 40 mg mL⁻¹ polystyrene beads (BioBeads SM2 Adsorbant, Biorad Laboratories) to the samples for 4 h. After incubation, BioBeads were separated from the GUVs by centrifugation (1000g, 10 min) and removed. The residual amount of DM after adsorption was not estimated. Empty control vesicles were prepared in a similar manner omitting the protein addition step. Electrophysiology recordings were made immediately following protein reconstitution.

Electrophysiology

All lipid bilayer experiments were performed using a planar patch clamp system (Port-a-Patch, Nanion Technologies GmbH), using borosilicate glass chip with an aperture diameter of approximately 1 µm. Channel activity was recorded in symmetrical condition comprised of 10 mM Na-Hepes, 200 mM NaCl, pH 7 (adjusted with NaOH). To determine the permeability of Na⁺ and K⁺, the reversal potential was measured in asymmetric conditions where the internal solution contained 10 mM Na-Hepes, 200 mM NaCl, pH 7 (adjusted with NaOH) and the external solution contained 10 mM Hepes, 110 mM KCl, pH 7 (adjusted with KOH). Data were filtered at 3 kHz (Bessel filter, HEKA amplifier, Lambrecht/Pfalz, Germany) digitized at a sampling rate of 50 kHz and analyzed with Clampfit (Axon Instruments). Bilayer formation process was computer controlled by PatchControl software (Nanion). The permeability ratio was estimated as in Ref. [32,40], according to equation $\frac{P_{Na}}{P_K} = \frac{a_{Na} [\exp(\frac{E_{rev} F}{RT})] [\exp(\frac{E_{rev} F}{RT}) + 1]}{4a_{se}}$. *R*, *T*, *F*, and *E_{rev}* are the gas constant, absolute temperature, Faraday constant, and the reversal potential, respectively. Activity coefficients for Na⁺ and K⁺ were estimated as follows: $a_s = \gamma_s [X_s]$, where activity, *a_s*, is the effective concentration of an ion in solution, *s* is related to the nominal concentration [*X_s*] by the activity coefficient, γ_s . γ_s was calculated from the Debye-Hückel equation: $\log_{10} \gamma_s = \frac{-0.51 \cdot z_s \sqrt{\mu}}{1 + 3.8 z_s \sqrt{\mu}}$ where μ is the ionic strength of the solution, *z_s* is the charge on the ion, and α_s is the effective diameter of the hydrated ion in nanometers.

Results and discussion

Based on the observation that membrane proteins can be successfully expressed in cell-free systems [10] and that *E. coli* is a suitable host for expressing the bacterial tetrameric pore-only Na_vSp1p fused to a maltose-binding protein and a hexa-His tag [32], we tested the expression of Na_vSp1p in a cell-free system derived from a bacterial extract S12 from *E. coli*. As Na_vSp1p is a membrane protein, we first undertook a detergent screen to effectively solubilize Na_vSp1p in CPFE. Based on literature research [41], we tried four non-polar detergents Brij®-58, Brij®-78, DDM and DM. The expression of Na_vSp1p was verified by immunoblotting in the crude expression mixture using anti-His antibodies (Fig. 1). We observed an equally strong signal both in the total reaction (T) and in the supernatant (S) indicating that the protein




Fig. 1. Western blot analysis of the solubilization Na_vSp1p expressed cell-free with four different detergents (Brij-78, DM, Brij-58, DDM). The blot is digitally overlaid with a picture of the same nitrocellulose membrane stained with Ponceau S ('S' denotes 'supernatant', 'T' denotes 'total reaction'). The frame marks the expected size of the fusion protein Na_vSp1p 62.1 kDa.




Fig. 2. Purification of Na_vSp1p. (A) Size-exclusion chromatogram, full and dashed line mark the absorbance at 280 nm of the Na_vSp1p obtained from cell-free expression and *E. coli* cells respectively. Asterisk indicates the collected peak (elution volume of the tetrameric Na_vSp1p (67.3 kDa) is 11.7–13.2 mL and 11.5–13.5 mL for cell-free expression and *E. coli* sample, respectively). We note that the large void volume in the sample expressed *in vitro* is likely due to excess DNA in the reaction mixture. (B, C) SDS-PAGE analyses of collected peaks stained with Coomassie. The asterisk marks the band for monomeric Na_vSp1p at 15.9 kDa from cell-free expression (B) and from *E. coli* cells (C).




Fig. 3. Corrected FTIR spectra of Na_vSp1p. The amide I (1600–1700 cm⁻¹) and amide II (~1550 cm⁻¹) vibrations of the polypeptide backbones are marked with one and two asterisks respectively.

was not only expressed, but also effectively solubilized in all detergents. Next, we optimized the incubation time and temperature. Changing the initial incubation time from 2 h to 4 h and overnight, did not result in a significant change in the yield. A temperature drop from 30 °C to 19 °C inhibited expression in the environment of DM and DDM and but did not change the expression in Brij-78 and Brij-58. We chose an incubation temperature and time of 30 °C and 2 h, respectively, and performed expression in DM. The protein was purified using affinity and size exclusion chromatography[32]. Optimization of the amount of protease for cleavage of the maltose binding domain at 8 °C yielded an optimized ratio of 1 mg: 0.28 mg (protein: protease).

We also expressed Na_vSp1p in *in vivo* in *E. coli* as described [32]. The size-exclusion chromatogram shows that Na_vSp1p from both expression systems has an identical elution volume (Fig. 2A). This is indicative of elution as a tetramer (67.3 kDa). SDS-PAGE analyses under denaturing conditions (Fig. 2B and C) showed that purification yields a pure protein in both cases and we observed a monomeric Na_vSp1p with the size of 15.9 kDa. The total yield was 20 μg purified protein per mL of cell-free reaction, which is sufficient for production of milligram amounts. For comparison, the yields of purified cell-free expressed human voltage-dependent anion channel-1 and bacteriorhodopsin were reported as 200–300 μg/mL and 24 μg/mL, respectively[11,13], both produced in batch mode format using *E. coli* extract.

To confirm that the ion channel was folded correctly, we recorded an FTIR spectrum (Fig. 3). The vibrational frequency of the amide I band (1656 cm⁻¹), and the peak shape of the amide I and II bands support the notion that the protein is alpha-helical [42].

Electrophysiological characterization of the Na_vSp1p, produced *in vitro*, reconstituted in planar lipid bilayers was analyzed on the single channel level. The single channel recordings showed that the




Fig. 4. Functional characterization of Na_vSp1p (A) Na_vSp1p reconstituted in planar lipid bilayer recorded in an asymmetric buffer solution, 110 mM KCl, 10 mM HEPES, pH 7 and 200 mM NaCl, 10 mM HEPES, pH 7 external and internal solution respectively, at different voltages as indicated. The closed channel current level is indicated. (B) Single channel IV relationships for Na_vSp1p channels in an asymmetric buffer solution, 110 mM KCl, 10 mM HEPES, pH 7 [K]_o and 200 mM NaCl, 10 mM HEPES, pH7 [Na]_i external and internal solution, respectively.

protein is a functional ion channel with a selectivity for sodium over potassium ions. We measured the conductance levels while varying the holding potential between -100 mV and $+100$ mV (Fig. 4A). The conductance of NavSp1 was 32.5 ± 2.8 pS under symmetrical condition (200 mM NaCl) which is in a good agreement with the previously reported value for the $\text{Na}_v\text{Sp1p}$ expressed in *E. coli* [32]. To further investigate the $\text{Na}_v\text{Sp1p}$, we measured its selectivity for sodium over potassium by measuring the reversal potential in a set of asymmetric ion conditions. The current – voltage relationship under asymmetric condition (110 mM KCl extracellular/200 mM NaCl intracellular) showed that $\text{Na}_v\text{Sp1p}$ has a preference for ions of $\text{Na}^+:\text{K}^+ = 1:0.27$ (Fig. 4B). This indicates that the expressed sodium channel is indeed functional and that it is selective for sodium over potassium ions.

In conclusion, we established a stable and functional cell-free expression system for $\text{Na}_v\text{Sp1p}$. The yield (0.02 mg purified protein/mL extract) is comparable to other membrane proteins [10,11,13]. The ion channel is a homo tetramer with functional characteristic resembling the protein expressed in *E. coli* [32]. This adds another functional membrane protein to the list of targets that can be produced by *in vitro* expression [10] and also to the list of *in vitro* expressed sodium channels, coming two decades after expression of a mechanosensitive renal Na_v [43] as the first from the family of bacterial voltage-gated Na_v s. The expression system can be adopted for efficient site-specific incorporation of isotope labeled amino acid, enabling the bio-spectroscopic investigation of this ion channel protein.

Acknowledgments

We thank Prof. Daniel Minor for the donation of the plasmids for $\text{Na}_v\text{Sp1p}$ and HRV 3C protease. This work was funded by grants from the Swedish Research Council, and the European Research Council through agreement number 279944.

References

[1] B. Hille, in: *Ion Channels Excit. Membr.*, Sinauer Associates, Inc., Sunderland, MA, 2001, pp. 1–22. http://pub.ist.ac.at/Pubs/courses/migrated/2012/introductionto Neuroscience1/docs/Lectures May 13,15/Hille_Ion Channels of Excitable Membranes_Chapter1.pdf (accessed March 16, 2014).

[2] W.A. Catterall, From ionic currents to molecular mechanisms, *Neuron* 26 (2000) 13–25. [http://dx.doi.org/10.1016/S0896-6273\(00\)81133-2](http://dx.doi.org/10.1016/S0896-6273(00)81133-2).

[3] W.A. Catterall, Voltage-dependent gating of sodium channels: correlating structure and function, *Trends Neurosci.* 9 (1986) 7–10. [http://dx.doi.org/10.1016/0166-2236\(86\)90004-4](http://dx.doi.org/10.1016/0166-2236(86)90004-4).

[4] D. Ren, B. Navarro, H. Xu, L. Yue, Q. Shi, D.E. Clapham, A prokaryotic voltage-gated sodium channel, *Science* 294 (2001) 2372–2375. <http://dx.doi.org/10.1126/science.1065635>.

[5] J. Payandeh, T. Scheuer, N. Zheng, W.A. Catterall, The crystal structure of a voltage-gated sodium channel, *Nature* 475 (2011) 353–358. <http://dx.doi.org/10.1038/nature10238>.

[6] E.C. McCusker, N. D'Avanzo, C.G. Nichols, B.A. Wallace, Simplified bacterial "pore" channel provides insight into the assembly, stability, and structure of sodium channels, *J. Biol. Chem.* 286 (2011) 16386–16391. <http://dx.doi.org/10.1074/jbc.C111.228122>.

[7] D.A. Doyle, J. Morais Cabral, R.A. Pfuetzner, A. Kuo, J.M. Gulbis, S.L. Cohen, The structure of the potassium channel: molecular basis of K^+ conduction and selectivity, *Science* 280 (1998) 69–77. <http://www.ncbi.nlm.nih.gov/pubmed/9525859> (accessed March 11, 2014).

[8] D.A. Köpfer, C. Song, T. Gruene, G.M. Sheldrick, U. Zachariae, B.L. de Groot, Ion permeation in K^+ channels occurs by direct Coulomb knock-on, *Science* 346 (2014) 352–355. <http://dx.doi.org/10.1126/science.1254840> (80–).

[9] S.Y. Noskov, S. Bernèche, B. Roux, Control of ion selectivity in potassium channels by electrostatic and dynamic properties of carbonyl ligands, *Nature* 431 (2004) 830–834. <http://dx.doi.org/10.1038/nature02943>.

[10] L. Isaksson, J. Enberg, R. Neutze, B. Göran Karlsson, A. Pedersen, Expression screening of membrane proteins with cell-free protein synthesis, *Protein Expr. Purif.* 82 (2012) 218–225. <http://dx.doi.org/10.1016/j.pep.2012.01.003>.

[11] S. Sonar, N. Patel, W. Fischer, K.J. Rothschild, Cell-free synthesis, functional refolding, and spectroscopic characterization of bacteriorhodopsin, an integral membrane protein, *Biochemistry* 32 (1993) 13777–13781.

[12] M.A. Goren, B.G. Fox, Wheat germ cell-free translation, purification, and assembly of a functional human stearoyl-CoA desaturase complex, *Protein Expr. Purif.* 62 (2008) 171–178. <http://dx.doi.org/10.1016/j.pep.2008.08.002>.

[13] A. Deniaud, L. Liguori, I. Blesneac, J.L. Lenormand, E. Pebay-Peyroula, Crystallization of the membrane protein hVDAC1 produced in cell-free system, *Biochim. Biophys. Acta* 1798 (2010) 1540–1546. <http://dx.doi.org/10.1016/j.bbame.2010.04.010>.

[14] F. Bernhard, Y. Tozawa, Cell-free expression—making a mark, *Curr. Opin. Struct. Biol.* 23 (2013) 374–380. <http://dx.doi.org/10.1016/j.sbi.2013.03.012>.

[15] C. Klammt, D. Schwarz, F. Löhr, B. Schneider, V. Dötsch, F. Bernhard, Cell-free expression as an emerging technique for the large scale production of integral membrane protein, *FEBS J.* 273 (2006) 4141–4153. <http://dx.doi.org/10.1111/j.1742-4658.2006.05432.x>.

[16] A.S. Spirin, High-throughput cell-free systems for synthesis of functionally active proteins, *Trends Biotechnol.* 22 (2004) 538–545. <http://dx.doi.org/10.1016/j.tibtech.2004.08.012>.

[17] B.W. Jarecki, S. Makino, E.T. Beebe, B.G. Fox, B. Chanda, Function of Shaker potassium channels produced by cell-free translation upon injection into *Xenopus* oocytes, *Sci. Rep.* 3 (2013) 1040. <http://dx.doi.org/10.1038/srep01040>.

[18] M.M. Falk, L.K. Buehler, N.M. Kumar, N.B. Gilula, Cell-free synthesis and assembly of connexins into functional gap junction membrane channels, *EMBO J.* 16 (1997) 2703–2716. <http://dx.doi.org/10.1093/emboj/16.10.2703>.

[19] L.K. Lyford, Cell-free expression and functional reconstitution of homo-oligomeric alpha 7 nicotinic acetylcholine receptors into planar lipid bilayers, *J. Biol. Chem.* 274 (1999) 25675–25681. <http://dx.doi.org/10.1074/jbc.274.36.25675>.

[20] L.T. Tegler, K. Corin, J. Hillger, B. Wassie, Y. Yu, S. Zhang, Cell-free expression, purification, and ligand-binding analysis of *Drosophila melanogaster* olfactory receptors DmOR67a, DmOR85b and DmORCO, *Sci. Rep.* 5 (2015) 7867. <http://dx.doi.org/10.1038/srep07867>.

[21] R.L. Rosenberg, J.E. East, Cell-free expression of functional Shaker potassium channels, *Nature* 360 (1992) 166–169. <http://dx.doi.org/10.1038/360166a0>.

[22] S.K. Dondapati, M. Kreir, R.B. Quast, D.A. Wüstenhagen, A. Brüggemann, N. Fertig, et al., Membrane assembly of the functional KcsA potassium channel in a vesicle-based eukaryotic cell-free translation system, *Biosens. Bioelectron.* 59 (2014) 174–183. <http://dx.doi.org/10.1016/j.bios.2014.03.004>.

[23] M.S. Awaysda, I.I. Ismailov, B.K. Berdiev, D.J. Benos, A cloned renal epithelial Na^+ channel protein displays stretch activation in planar lipid bilayers, *Am. J. Physiol.* 268 (1995) C1450–C1459. <http://www.ncbi.nlm.nih.gov/pubmed/7611365> (accessed February 23, 2015).

[24] T. Kigawa, T. Yabuki, Y. Yoshida, M. Tsutsui, Y. Ito, T. Shibata, et al., Cell-free production and stable-isotope labeling of milligram quantities of proteins, *FEBS Lett.* 442 (1999) 15–19. <http://www.ncbi.nlm.nih.gov/pubmed/9923595> (accessed December 28, 2014).

[25] C.R. Baiz, M. Reppert, A. Tokmakoff, Introduction to protein 2D IR Spectroscopy, (2012) 1–38.

[26] S. Woutersen, Y. Mu, G. Stock, P. Hamm, Hydrogen-bond lifetime measured by time-resolved 2D-IR spectroscopy: N-methylacetamide in methanol, *Chem. Phys.* 266 (2001) 137–147. [http://dx.doi.org/10.1016/S0301-0104\(01\)00224-5](http://dx.doi.org/10.1016/S0301-0104(01)00224-5).

[27] Z. Ganim, A. Tokmakoff, A. Vaziri, Vibrational excitons in ionophores: experimental probes for quantum coherence-assisted ion transport and selectivity in ion channels, *New J. Phys.* 13 (2011) 113030. <http://dx.doi.org/10.1088/1367-2630/13/11/113030>.

[28] C. Fang, R.M. Hochstrasser, Two-dimensional infrared spectra of the 13C=18O isotopomers of alanine residues in an alpha-helix, *J. Phys. Chem. B* 109 (2005) 18652–18663. <http://dx.doi.org/10.1021/jp052525p>.

[29] S. Ye, E. Zaitseva, G. Caltabiano, G.F.X. Schertler, T.P. Sakmar, X. Deupi, et al., Tracking G-protein-coupled receptor activation using genetically encoded infrared probes, *Nature* 464 (2010) 1386–1389. <http://dx.doi.org/10.1038/nature08948>.

[30] C.J. Noren, S.J. Anthony-Cahill, M.C. Griffith, P.G. Schultz, A general method for site-specific incorporation of unnatural amino acids into proteins, *Science* 244 (1989) 182–188. <http://www.ncbi.nlm.nih.gov/pubmed/2649980> (accessed October 28, 2014).

[31] K.V. Loscha, A.J. Herlt, R. Qi, T. Huber, K. Ozawa, G. Otting, Multiple-site labeling of proteins with unnatural amino acids, *Angew. Chem. Int. Ed. Engl.* 51 (2012) 2243–2246. <http://dx.doi.org/10.1002/anie.201108275>.

[32] D. Shaya, M. Kreir, R.A. Robbins, S. Wong, J. Hammon, A. Brüggemann, et al., Voltage-gated sodium channel (Nav) protein dissection creates a set of functional pore-only proteins, *Proc. Natl. Acad. Sci. U.S.A.* 108 (2011) 12313–12318. <http://dx.doi.org/10.1073/pnas.1106811108>.

[33] A. Pedersen, K. Hellberg, J. Enberg, B.G. Karlsson, Rational improvement of cell-free protein synthesis, *N. Biotechnol.* 28 (2011) 218–224. <http://www.sciencedirect.com/science/article/pii/S1871678410004917> (accessed January 20, 2014).

[34] S. Peuker, H. Andersson, K.S. Maiti, S. Niebling, A. Pedersen, M. Erdelyi et al., Isotope-edited proteins for site-directed infrared spectroscopy, Submitted (n.d.).

[35] J. Grodberg, J.J. Dunn, OmpT encodes the *Escherichia coli* outer membrane protease that cleaves T7 RNA polymerase during purification, *J. Bacteriol.* 170 (1988) 1245–1253.

[36] P. Davanloo, A.H. Rosenberg, J.J. Dunn, F.W. Studier, Cloning and expression of the gene for bacteriophage T7 RNA polymerase, *Proc. Natl. Acad. Sci. U.S.A.* 81 (1984) 2035–2039. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=345431&tool=pmcentrez&rendertype=abstract> (accessed December 16, 2014).

- 513 [37] M. Kreir, C. Farre, M. Beckler, M. George, N. Fertig, Rapid screening of
514 membrane protein activity: electrophysiological analysis of OmpF
515 reconstituted in proteoliposomes, *Lab Chip* 8 (2008) 587–595, [http://](http://dx.doi.org/10.1039/b713982a)
516 dx.doi.org/10.1039/b713982a.
- 517 [38] O. Gassmann, M. Kreir, C. Ambrosi, J. Pranskevich, A. Oshima, C. Röling, et al.,
518 The M34A mutant of Connexin26 reveals active conductance states in pore-
519 suspending membranes, *J. Struct. Biol.* 168 (2009) 168–176, [http://dx.doi.org/](http://dx.doi.org/10.1016/j.jsb.2009.02.004)
520 [10.1016/j.jsb.2009.02.004](http://dx.doi.org/10.1016/j.jsb.2009.02.004).
- 521 [39] N. D'Avanzo, E.C. McCusker, A.M. Powl, A.J. Miles, C.G. Nichols, B.A. Wallace,
522 Differential lipid dependence of the function of bacterial sodium channels,
523 *PLoS ONE* 8 (2013) e61216, <http://dx.doi.org/10.1371/journal.pone.0061216>.
- [40] L. Yue, B. Navarro, D. Ren, A. Ramos, D.E. Clapham, The cation selectivity filter
of the bacterial sodium channel, *NaChBac*, *J. Gen. Physiol.* 120 (2002) 845–853.
[http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2229573&tool=](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2229573&tool=pmcentrez&rendertype=abstract)
pmcentrez&rendertype=abstract (accessed December 07, 2014).
- [41] D. Schwarz, C. Klammt, A. Koglin, F. Löhr, B. Schneider, V. Dötsch, et al.,
Preparative scale cell-free expression systems: new tools for the large scale
preparation of integral membrane proteins for functional and structural studies,
Methods 41 (2007) 355–369, <http://dx.doi.org/10.1016/j.ymeth.2006.07.001>.
- [42] F. Siebert, P. Hildebrandt, *Vibrational Spectroscopy in Life Science*, 2008.
- [43] I. Ismailov, K. Berdiev, D.J. Benos, A cloned renal epithelial Na⁺ channel protein
displays stretch activation in planar lipid bilayers, (1995).

524
525
526
527
528
529
530
531
532
533
534
535

UNCORRECTED PROOF